JZ Mock Set D Paper 1 Solutions

Recommended time: 2 hours

Calculators: not permitted

Format: 20 multiple-choice questions

This is a TMUA-style mock paper modelled on the Test of Mathematics for University Admission. The TMUA is used in admissions for mathematics, economics, computer science, and engineering courses at universities including Cambridge, Oxford, Imperial College London, UCL, LSE, Warwick, and Durham.

jzmaths.com

Paper Guidance

- These papers contain only hard, very hard, and occasionally *unreasonably hard* TMUA-style questions :). I have deliberately left out the routine and easier material, so please do not be discouraged if you find a paper difficult or need longer than the recommended time. That is exactly how it is meant to feel. Take your time, enjoy the challenge, and remember that struggling with difficult problems is one of the best ways to get better.
- As a rough and deliberately conservative guide, a student who scores above 27 out of 40 across one full set, meaning both papers in four hours, is very likely already working at around the 7.0+ level on the current TMUA scale. However, this is only intended as a guide, not something to worry about too much. The main purpose of these papers is improvement rather than measurement: meet difficult problems, struggle with them, learn from them, and focus on getting better each time.

Best of luck, and enjoy the challenge!

Last updated: 30 September 2026 at 21:38

Spotted an error? Please email jzmaths@hotmail.com.

Question 1

Find the coefficient of x^4 in the expansion of

$$\left(2 - x + \frac{x^2}{2}\right)^5.$$

A 50

B 60

C 70

D 80

E 90

Solution 1

Answer: E

One of a few standard questions. I may decide to replace it with a **non-standard** question at a later date, so take it easy for now!

jzmaths.com

Using the binomial expansion,

$$\left((2-x) + \frac{x^2}{2}\right)^5 = (2-x)^5 + 5(2-x)^4\left(\frac{x^2}{2}\right) + 10(2-x)^3\left(\frac{x^2}{2}\right)^2 + \cdots.$$

Only these three terms can contribute to the coefficient of x^4. Their contributions are

$$\begin{aligned}(2-x)^5 &: \binom{5}{4}2(-x)^4 = 10x^4,\\ 5(2-x)^4\left(\frac{x^2}{2}\right) &: 5\binom{4}{2}2^2(-x)^2\left(\frac{x^2}{2}\right) = 60x^4,\\ 10(2-x)^3\left(\frac{x^2}{2}\right)^2 &: 10\cdot 2^3\left(\frac{x^4}{4}\right) = 20x^4.\end{aligned}$$

Therefore, the coefficient of x^4 is

$$10 + 60 + 20 = 90.$$

Question 2

How many solutions are there to the equation

$$|\tan x| + \frac{1}{\tan x} = 3 \quad \text{for} \quad 0^\circ \leq x \leq 360^\circ?$$

A 2

B 4

C 5

D 6

E 8

Solution 2

Answer: D

jzmaths.com

Let $t = \tan x$, where $t \neq 0$. The equation becomes

$$|t| + \frac{1}{t} = 3.$$

If $t > 0$, then $t + \frac{1}{t} = 3$, so $t^2 - 3t + 1 = 0$. This quadratic has two positive roots, you can see this by sketching suitable graphs, such as $y = 1$ and $y = 3t - t^2$, there is no need to solve the equation for t. Each root gives two values of x for $0^\circ \leq x \leq 360^\circ$. Hence there are 4 solutions.

If $t < 0$, then $-t + \frac{1}{t} = 3$, so $t^2 + 3t - 1 = 0$. This quadratic has one negative root, which gives two values of x in the given interval. Hence there are 2 more solutions.

Therefore, the total number of solutions is $4 + 2 = 6$.

Question 3

The functions f and g are defined for all real x by

$$f(x) = |x - 2| + |x + 2|$$

and

$$g(x) = x^2 - 2x + 4.$$

The function h is defined by

$$h(x) = g(f(x)).$$

What is the minimum value of $h(x)$?

A 10

B 11

C 12

D 13

E 14

F 15

jzmaths.com

Solution 3

Answer: C

Since $f(x) = |x - 2| + |x + 2|$ is the sum of the distances from x to 2 and -2, we have $f(x) \geq 4$, with equality for $-2 \leq x \leq 2$.

Also,

$$g(t) = t^2 - 2t + 4 = (t - 1)^2 + 3.$$

For $t \geq 4$, $g(t)$ is increasing. Therefore, $g(f(x))$ is minimised when $f(x) = 4$.

Hence the minimum value is $g(4) = 16 - 8 + 4 = 12$.

Question 4

Find the total length of interval(s) on which

$$f(x) = \frac{2}{5}x^{5/2} - 2x^{3/2} + 4x^{1/2} \qquad x > 0$$

is decreasing.

A $\frac{1}{2}$

B 1

C 2

D 3

E 4

F $\frac{5}{2}$

jzmaths.com

Solution 4

Answer: B

Differentiate:

$$f'(x) = x^{3/2} - 3x^{1/2} + 2x^{-1/2}.$$

Factorise:

$$f'(x) = x^{-1/2}(x^2 - 3x + 2) = x^{-1/2}(x-1)(x-2).$$

Since $x > 0$, we have $x^{-1/2} > 0$, so the sign of $f'(x)$ is the sign of $(x-1)(x-2)$.

The function is decreasing when $f'(x) \leq 0$, so

$$(x-1)(x-2) \leq 0.$$

Hence $1 \leq x \leq 2$. Therefore the total length of the interval on which f is decreasing is $2 - 1 = 1$.

Question 5

Find the area of the region in the (x, y)-plane consisting of all points satisfying both

$$|x| + |y| \leq 2 \quad \text{and} \quad |x - 1| + |y| \leq 2.$$

A 3

B $\frac{9}{4}$

C $\frac{7}{2}$

D 4

E $\frac{9}{2}$

F 6

G $\frac{15}{2}$

H 8

jzmaths.com

Solution 5

Answer: E

The inequality $|x| + |y| \leq 2$ describes a diamond centred at $(0, 0)$, while $|x - 1| + |y| \leq 2$ describes an identical diamond centred at $(1, 0)$.

Sketch out the two diamonds, then you will see that their common region is also a diamond. Its leftmost and rightmost vertices are $(-1, 0)$ and $(2, 0)$, so its horizontal diagonal has length 3.

Its upper and lower vertices are

$$\left(\frac{1}{2}, \frac{3}{2}\right) \quad \text{and} \quad \left(\frac{1}{2}, -\frac{3}{2}\right),$$

so its vertical diagonal also has length 3.

Therefore, the area of the common region is

$$\frac{1}{2} \cdot 3 \cdot 3 = \frac{9}{2}.$$

Question 6

An arithmetic progression has $(2n+1)$ terms, where n is a positive integer. Let S_o denote the sum of the terms in odd-numbered positions (the 1st, 3rd, 5th, ... terms) and let S_e denote the sum of the terms in even-numbered positions (the 2nd, 4th, 6th, ... terms).

It is given that

$$S_o + S_e = 505 \qquad \text{and} \qquad S_o - S_e = 5.$$

What is the value of n?

A 40

B 50

C 51

D 55

E 56

jzmaths.com

Solution 6

Answer: B

Since there are an odd number $2n+1$ terms, there must be a middle term, call it M, then $S_o = (n+1)M$ and $S_e = nM$. Therefore

$$S_o + S_e = (n+1)M + nM = (2n+1)M,$$
$$S_o - S_e = (n+1)M - nM = M.$$

From $S_o - S_e = 5$ we get $M = 5$. Substituting into $S_o + S_e = 505$ gives $(2n+1) \cdot 5 = 505$, so $2n+1 = 101$ and hence $n = 50$.

Question 7

Find the value of

$$\sum_{r=1}^{100} \frac{1}{\sqrt{r}+\sqrt{r-1}}.$$

A 0

B 1

C 9

D 10

E 11

F 20

Solution 7

Answer: D

jzmaths.com

Rationalising each term gives

$$\frac{1}{\sqrt{r}+\sqrt{r-1}} = \frac{1}{\sqrt{r}+\sqrt{r-1}} \times \frac{\sqrt{r}-\sqrt{r-1}}{\sqrt{r}-\sqrt{r-1}} = \sqrt{r}-\sqrt{r-1}.$$

Therefore,

$$\begin{aligned}\sum_{r=1}^{100} \frac{1}{\sqrt{r}+\sqrt{r-1}} &= \sum_{r=1}^{100} \left(\sqrt{r}-\sqrt{r-1}\right) \\ &= \sqrt{100}-\sqrt{0} \\ &= 10.\end{aligned}$$

Question 8

The two curves C_1 and C_2 in the xy-plane have equations

$$x^2 + y^2 - 2y - 12 = 0 \text{ and } x^2 + y^2 - 2x - 4y = 0.$$

Let P be a point on C_1 and Q be a point on C_2. What is the **least** possible value of the length of PQ?

A $\sqrt{2} + \sqrt{5} - \sqrt{13}$

B $\sqrt{13} - \sqrt{5} - \sqrt{2}$

C $\sqrt{13} - \sqrt{2}$

D 0

E $\sqrt{13} + \sqrt{5} - \sqrt{2}$

Solution 8

Answer: D

C_1: $x^2 + (y-1)^2 = 13$, so its centre is $A = (0, 1)$ and its radius is $\sqrt{13}$.

C_2: $(x-1)^2 + (y-2)^2 = 5$, so its centre is $B = (1, 2)$ and its radius is $\sqrt{5}$.

Now $AB = \sqrt{2}$. The circles intersect if $\sqrt{2} + \sqrt{5} > \sqrt{13}$. Since

$$(\sqrt{2} + \sqrt{5})^2 = 7 + 2\sqrt{10} > 13,$$

the circles intersect.

Hence we may choose $P = Q$ at an intersection point, so the least possible value of PQ is 0. Indeed the whole point of this question is to get you to determine how close they really are.

Alternatively, shortcut to mushrooms: subtracting the two equations gives $2x + 2y - 12 = 0$, so $y = 6 - x$. Substituting this into either equation gives

$$x^2 - 5x + 6 = 0,$$

which has two distinct real roots. Hence the circles intersect, so the least possible value of PQ is 0.

jzmaths.com

Question 9

Let x be any real number, find the minimum value of

$$4^{\sin^2 x} - 5 \cdot 2^{\sin^2 x} + 7.$$

A There is no minimum value.

B 1

C $\frac{5}{4}$

D 3

E 7

F $\frac{3}{4}$

Solution 9

Answer: B

jzmaths.com

Another completing the square question with a twist!

Let $u = 2^{\sin^2 x}$. Since $\sin^2 x$ takes every value in $[0, 1]$ (and only those values) as x varies over $\mathbb{R}$, u ranges over $[2^0, 2^1] = [1, 2]$.

Note that $4^{\sin^2 x} = (2^{\sin^2 x})^2 = u^2$, so the expression becomes

$$g(u) = u^2 - 5u + 7 = \left(u - \frac{5}{2}\right)^2 + \frac{3}{4}.$$

The minimum of g is $\frac{3}{4}$ if u is allowed to be $\frac{5}{2}$, it is not! As u varies between 1 to 2, given the shape of $g(u)$, the minimum is clearly at $u = 2$:

$$g(2) = 4 - 10 + 7 = 1.$$

Question 10

Given that $x^2 - 4x + 1$ is a factor of

$$2x^4 - 11x^3 + bx^2 + cx + 6,$$

what is the value of $b + c$?

A -47

B -27

C -7

D 13

E 47

F 53

jzmaths.com

Solution 10

Answer: C

Since $x^2 - 4x + 1$ is a factor and the quartic has leading coefficient 2 and constant term 6, write

$$2x^4 - 11x^3 + bx^2 + cx + 6 = (x^2 - 4x + 1)(2x^2 + px + q).$$

Expanding the right-hand side gives

$$2x^4 + (p - 8)x^3 + (q - 4p + 2)x^2 + (p - 4q)x + q.$$

Comparing coefficients:

$$\begin{aligned} x^0: &\quad q = 6, \\ x^3: &\quad p - 8 = -11 \implies p = -3, \\ x^2: &\quad b = q - 4p + 2 = 6 + 12 + 2 = 20, \\ x^1: &\quad c = p - 4q = -3 - 24 = -27. \end{aligned}$$

Therefore $b + c = 20 + (-27) = -7$.

Remark: The roots of $x^2 - 4x + 1$ are $x = 2 \pm \sqrt{3}$, so direct substitution would not be wise!

Question 11

Let $f(x) = -x^3 + kx^2 + 3$, where k is a real constant. Suppose that $f(x) = 0$ has exactly 3 distinct real roots.

Which of the following equations must also have exactly 3 distinct real roots?

I $\quad -f(x) + a = 0$, where a is an arbitrary real constant

II $\quad 5 - 2f(2 - 3x) = 0$

III $\quad f(-|x|) - 3 = 0$

A I only

B II only

C III only

D I and III only

E I and II only

F all of them

G II and III only

H none of them

jzmaths.com

Solution 11

Answer: G

We have $f'(x) = x(2k - 3x)$, so the stationary points are at $x = 0$ and $x = \frac{2k}{3}$.

Since $f(0) = 3 > 0$ and $f(x) = 0$ has three distinct real roots, $x = 0$ must be the local maximum and the other stationary point must be a local minimum below the x-axis. Hence $\frac{2k}{3} < 0$, so $k < 0$.

Equation I need not have three distinct real roots, since a can be chosen sufficiently large that both stationary values of $-f(x) + a$ are positive, giving only one real root.

For II, let $u = 2 - 3x$. Then

$$5 - 2f(2 - 3x) = 0 \quad \iff \quad f(u) = \frac{5}{2}.$$

The local maximum value is 3 and the local minimum value is negative. Since $0 < \frac{5}{2} < 3$, the line $y = \frac{5}{2}$ intersects the graph three times. As $u = 2 - 3x$ is one-to-one, II has three distinct real roots.

For III,

$$f(-|x|) - 3 = |x|^3 + kx^2 = x^2(|x| + k).$$

Since $k < 0$, the equation has the three distinct roots $x = 0$, $x = k$ and $x = -k$.

Therefore, II and III only must have exactly three distinct real roots.

jzmaths.com

Question 12

The curve $x = y^2 - 8y + 19$ is rotated 90° clockwise about $P_1 = (1, 2)$, and then 90° clockwise about P_2. The final curve is $x = -y^2 + 2y - 7$. Given P_2 lies on the line $y = 5$, find the x-coordinate of P_2.

A -1

B -2

C -3

D 0

E 1

F 2

G 3

H 4

jzmaths.com

Solution 12

Answer: A

Let $P_2 = (a, 5)$.

First rewrite the original curve:

$$x = y^2 - 8y + 19 = (y - 4)^2 + 3.$$

So its vertex is $(3, 4)$.

The final curve is

$$x = -y^2 + 2y - 7 = -(y - 1)^2 - 6.$$

So its vertex is $(-6, 1)$.

A rotation maps the vertex of the parabola to the vertex of the new parabola, so track the vertex only.

The key observation here is: a 90° clockwise rotation about (h, k) sends a point (x, y) to $(h + (y - k), k - (x - h)) = (h + y - k, k - x + h)$, which can be seen from a diagram (check it out yourself).

Rotating $(3, 4)$ about $P_1 = (1, 2)$ gives

$$(1 + 4 - 2, 2 - 3 + 1) = (3, 0).$$

Now rotate $(3, 0)$ about $P_2 = (a, 5)$:

$$(a + 0 - 5, 5 - 3 + a) = (a - 5, a + 2).$$

This must equal the final vertex $(-6, 1)$, so

$$a - 5 = -6$$

and hence $a = -1$.

Therefore the x-coordinate of P_2 is -1.

jzmaths.com

Question 13

The graph of $y = \sin x$ is transformed by the following five operations, applied in the order given:

I a reflection in the line $y = -3$;

II a stretch by scale factor 2 parallel to the x-axis;

III a translation of $\frac{\pi}{3}$ in the negative x-direction.

IV a reflection in the line $x = 0$;

V a reflection in the line $x = \frac{4\pi}{3}$.

Which of the following is the equation of the resulting curve?

A

$$y = -3 + \sin\left(\frac{x}{2} + \frac{5\pi}{12}\right)$$

B

$$y = -6 - \sin\left(\frac{x}{2} - \frac{5\pi}{12}\right)$$

C

$$y = -6 + \sin\left(\frac{x}{2} - \frac{\pi}{6}\right)$$

D

$$y = -3 + \sin\left(\frac{x}{2} - \frac{5\pi}{12}\right)$$

E

$$y = 6 + \sin\left(\frac{x}{2} - \frac{5\pi}{12}\right)$$

jzmaths.com

Solution 13

Answer: C

The reflection in the line $y = -3$ affects only the y-coordinate, while transformations II to V affect only the x-coordinate. We may therefore apply transformation I last.

Starting with $y = \sin x$, the stretch and translation give

$$y = \sin\left(\frac{x}{2} + \frac{\pi}{6}\right).$$

Reflecting in $x = 0$ gives

$$y = \sin\left(-\frac{x}{2} + \frac{\pi}{6}\right).$$

The lines of symmetry of this curve satisfy

$$-\frac{x}{2}+\frac{\pi}{6}=\frac{\pi}{2}+n\pi,$$

so $x=-\frac{2\pi}{3}-2n\pi$. The least positive value is $\frac{4\pi}{3}$, so reflection in $x=\frac{4\pi}{3}$ leaves the curve unchanged.

Finally, reflection in $y=-3$ changes y to $-6-y$, giving

$$y=-6-\sin\left(-\frac{x}{2}+\frac{\pi}{6}\right)=-6+\sin\left(\frac{x}{2}-\frac{\pi}{6}\right).$$

jzmaths.com

Question 14

The equation

$$\cos x + \cos^3 x + \cos^5 x + \cos^7 x + \ldots = \tan x$$

has how many solutions for $0 \leq x < 2\pi$?

A 0

B 1

C 2

D 3

E 4

F infinitely many

Solution 14

Answer: C

jzmaths.com

The left-hand side is a geometric series with first term $\cos x$ and common ratio $\cos^2 x$. It converges for $\cos^2 x < 1$, so we must exclude $x = 0$ and $x = \pi$, where $\cos^2 x = 1$. The right-hand side $\tan x$ is not defined for $x = \frac{\pi}{2}$ and $x = \frac{3\pi}{2}$, so we also exclude these x.

When the series converges, its sum to infinity is

$$\frac{\cos x}{1 - \cos^2 x} = \frac{\cos x}{\sin^2 x}.$$

Substituting into the equation, we get

$$\frac{\cos x}{\sin^2 x} = \tan x \quad \Longleftrightarrow \quad \frac{\cos x}{\sin^2 x} = \frac{\sin x}{\cos x}.$$

Multiplying both sides by $\sin^2 x \cdot \cos x$, which is nonzero as we precisely excluded the values of x for which they are zero, simplify, eventually gives

$$\sin^3 x + \sin^2 x - 1 = 0$$

Let $u = \sin x$. Since $-1 \leq u \leq 1$, we need to count the solutions of $u^3 = 1 - u^2$ in this interval. For $-1 \leq u \leq 0$, we have $u^3 \leq 0$ and $1 - u^2 \geq 0$, so there are no solutions. For $0 \leq u \leq 1$, the

function u^3 is strictly increasing from 0 to 1, while $1 - u^2$ is strictly decreasing from 1 to 0. Hence they intersect exactly once, at some $u = k$ with $0 < k < 1$. The equation $\sin x = k$ has exactly two solutions for $0 \leq x < 2\pi$, so the original equation has 2 solutions.

jzmaths.com

Question 15

The circle C is inscribed in the triangle bounded by the three lines

$$y = 0, \qquad y = \sqrt{3}x, \qquad x + y = \sqrt{1} + \sqrt{2} + \sqrt{3}.$$

You may find $\tan 22.5^\circ = \sqrt{2} - 1$ useful.

Which of the following points lies inside C?

A $\left(\frac{1}{2}, 1\right)$

B $(2, 2)$

C $\left(\frac{3}{2}, \frac{5}{2}\right)$

D $\left(1, \frac{5}{3}\right)$

E $\left(3, \frac{3}{4}\right)$

Solution 15

Answer: D

jzmaths.com

Remark 1: This is a modified version of the question. Rob Drew, while making a walkthrough, spotted a wonderful unintended shortcut hidden in the options, thank you Rob! I have now diligently closed that particular escape route, so the rest of you can enjoy the full richness of the mathematics involved. :)

The key idea is that the centre of an inscribed circle lies at the intersection of the internal angle bisectors (GCSE-ish knowledge!). Since the radius to $y = 0$ is vertical, the y-coordinate of the centre is therefore equal to the radius. Once we have found the centre, we can simply calculate the distance from each given point to the centre to determine which one lies inside C.

At the origin, the angle between $y = 0$ and $y = \sqrt{3}x$ is 60°, so the angle bisector makes an angle of 30° with the positive x-axis. Therefore its equation is

$$y = x \tan 30^\circ = \frac{x}{\sqrt{3}}.$$

The line $x + y = 1 + \sqrt{2} + \sqrt{3}$ meets the x-axis at

$$(1 + \sqrt{2} + \sqrt{3}, 0).$$

At this point, the internal angle is between the negative x-axis and the line $x + y = 1 + \sqrt{2} + \sqrt{3}$. Since the line has gradient -1, the angle bisector makes an angle of $22.5°$ above the negative x-axis.

Using $\tan 22.5° = \sqrt{2} - 1$, the second angle bisector is

$$y = (\sqrt{2} - 1)(1 + \sqrt{2} + \sqrt{3} - x).$$

Now intersect the two angle bisectors:

$$\frac{x}{\sqrt{3}} = (\sqrt{2} - 1)(1 + \sqrt{2} + \sqrt{3} - x).$$

At this point, some of you may be able to spot the solution by inspection: $x = \sqrt{3}$. This makes the left-hand side equal to 1, while the right-hand side becomes a difference of two squares, $(\sqrt{2} - 1)(\sqrt{2} + 1) = 1$.

If you do not spot this, simply solve the one-variable equation normally to obtain $x = \sqrt{3}$. Then, substituting back, we quickly deduce $y = 1$.

Therefore the centre is $(\sqrt{3}, 1)$. Since the circle touches the line $y = 0$, its radius is therefore 1.

Now we test each of the points. A point is inside the circle if and only if its distance from the centre is less than 1.

For $\left(\frac{1}{2}, 1\right)$, the horizontal distance from the centre alone is $\sqrt{3} - \frac{1}{2} > 1$, so it is outside.

For $(2, 2)$, the vertical distance is already 1, and the horizontal distance is non-zero, so it is outside.

For $\left(\frac{3}{2}, \frac{5}{2}\right)$, the vertical distance from the centre is $\frac{3}{2} > 1$, so it is outside.

For $\left(1, \frac{5}{3}\right)$, the square of the distance from the centre is

$$(1 - \sqrt{3})^2 + \left(\frac{5}{3} - 1\right)^2 = \frac{40}{9} - 2\sqrt{3}.$$

Ouch. Is that less than 1? Probably, but rather than wrestling with it, let's check the final option first. :)

For $\left(3, \frac{3}{4}\right)$, the horizontal distance from the centre alone is $3 - \sqrt{3} > 1$, so it is outside.

Therefore, by elimination, the one option I deliberately did not bother to verify must be the correct one. Hence the point lying inside C is $\left(1, \frac{5}{3}\right)$.

For completeness, $\frac{40}{9} - 2\sqrt{3} < 1$ is indeed true. This is equivalent to $31 < 18\sqrt{3}$, and squaring gives $961 < 972$. So the point really does lie inside C.

Remark 2: To save myself from the further mild embarrassment of someone finding another very simple unintended shortcut, I set about looking for alternative methods myself. There is indeed

jzmaths.com

another method, but thankfully this one is a genuinely nice alternative solution rather than an escape route!

Here it is:

Let the radius be r. Since the centre lies on the 30° angle bisector and its y-coordinate is r, the centre must be $(\sqrt{3}r, r)$.

Now use the fact that the circle is also tangent to $x + y = 1 + \sqrt{2} + \sqrt{3}$. The perpendicular distance from $(\sqrt{3}r, r)$ to this line can be found to be

$$\frac{1 + \sqrt{2} + \sqrt{3} - (\sqrt{3} + 1)r}{\sqrt{2}}.$$

Since this distance is also the radius,

$$\frac{1 + \sqrt{2} + \sqrt{3} - (\sqrt{3} + 1)r}{\sqrt{2}} = r.$$

Hence

$$1 + \sqrt{2} + \sqrt{3} = (1 + \sqrt{2} + \sqrt{3})r,$$

so immediately $r = 1$, and therefore the centre is $(\sqrt{3}, 1)$.

Of course, there is still no guarantee that some enterprising student will not find another escape route I have completely missed. :)

jzmaths.com

Question 16

Let f be a real-valued function such that

$$\int_0^1 f(x)\,dx = 1$$

and $f(x) = f(x+1)$ for all real x.

Find

$$\sum_{r=1}^{100} \int_0^r f(r^r x)\,dx.$$

A 50

B 100

C 101

D 5050

E 2050

F 4950

jzmaths.com

Solution 16

Answer: D

Remark: The functions $f(x), f(4x), f(27x), f(256x), \ldots$ do not look especially friendly. You may wonder how much the coefficients of x inside f matter. It turns out that they do not matter at all!

For any positive integer k, the graph of $y = f(kx)$ is obtained from the graph of $y = f(x)$ by a horizontal stretch with scale factor $\frac{1}{k}$.

Since f has period 1, the function $f(kx)$ completes k periods on the interval $[0, 1]$. Each compressed period has integral

$$\frac{1}{k}\int_0^1 f(x)\,dx = \frac{1}{k}.$$

Therefore,

$$\int_0^1 f(kx)\,dx = k \cdot \frac{1}{k} = 1.$$

For each r, the number r^r is a positive integer, so

$$\int_0^1 f(r^r x)\,dx = 1.$$

Also, since r^r is an integer,

$$f(r^r(x+1)) = f(r^r x + r^r) = f(r^r x),$$

so $f(r^r x)$ has period 1.

Hence the interval $[0, r]$ consists of r intervals of length 1, each contributing an integral of 1. Therefore,

$$\int_0^r f(r^r x)\,dx = r.$$

Thus the required sum is

$$\sum_{r=1}^{100} r = 1 + 2 + 3 + \cdots + 100 = \frac{100 \cdot 101}{2} = 5050.$$

jzmaths.com

Question 17

A real number x satisfies

$$x^2 - 3x + 1 = 0.$$

For each non-negative integer n, define

$$p_n = x^n + \frac{1}{x^n}.$$

Find the value of p_6.

You may find it helpful to consider a recurrence relation of the form $p_n = 3p_{n-1} + kp_{n-2}$ for some constant k.

A 322

B 123

C 312

D 151

E 276

F 258

jzmaths.com

Solution 17

Answer: A

Remark: If a question says, **You may find it helpful**, this is often a euphemism for, **If you do not use this, you are unlikely to get very far!** So, although it may sound optional, you would be wise to follow the hint.

Since $x^2 - 3x + 1 = 0$ and $x \neq 0$, dividing by x gives

$$x + \frac{1}{x} = 3.$$

For $n \geq 2$,

$$\begin{aligned} 3p_{n-1} &= \left(x + \frac{1}{x}\right)\left(x^{n-1} + \frac{1}{x^{n-1}}\right) \\ &= x^n + x^{n-2} + \frac{1}{x^{n-2}} + \frac{1}{x^n} \\ &= p_n + p_{n-2}. \end{aligned}$$

Therefore,

$$p_n = 3p_{n-1} - p_{n-2},$$

so $k = -1$.

The initial values are $p_0 = 2$ and $p_1 = x + \frac{1}{x} = 3$.

We can now apply the recurrence repeatedly:

$$\begin{aligned}
p_2 &= 3p_1 - p_0 = 3(3) - 2 = 7,\\
p_3 &= 3p_2 - p_1 = 3(7) - 3 = 18,\\
p_4 &= 3p_3 - p_2 = 3(18) - 7 = 47,\\
p_5 &= 3p_4 - p_3 = 3(47) - 18 = 123,\\
p_6 &= 3p_5 - p_4 = 3(123) - 47 = 322.
\end{aligned}$$

Hence

$$p_6 = 322.$$

jzmaths.com

Question 18

Let $\lfloor x \rfloor$ denote the greatest integer less than or equal to x.

Find the value of

$$\int_{100}^{1} \log_{10}\left(\frac{\lfloor 2x \rfloor}{\lfloor 2x \rfloor + 1}\right) dx.$$

A -1

B $\log_{10} 99$

C $\log_{10} 101$

D $-\log_{10} 99$

E $-\log_{10} 101$

F 1

Solution 18

Answer: F

Remark: Students often find integrals involving step functions difficult, so here is an opportunity to practise! I have added a couple of extra twists to this one. Do you like them?!

Reversing the limits changes the sign of the integral:

$$\int_{100}^{1} \log_{10}\left(\frac{\lfloor 2x \rfloor}{\lfloor 2x \rfloor + 1}\right) dx = -\int_{1}^{100} \log_{10}\left(\frac{\lfloor 2x \rfloor}{\lfloor 2x \rfloor + 1}\right) dx.$$

For $\frac{k}{2} \leq x < \frac{k+1}{2}$, we have $\lfloor 2x \rfloor = k$. From $x = 1$ to $x = 100$, the values of $\lfloor 2x \rfloor$ are $2, 3, \ldots, 199$, each lasting for an interval of length $\frac{1}{2}$. Therefore

$$\int_{1}^{100} \log_{10}\left(\frac{\lfloor 2x \rfloor}{\lfloor 2x \rfloor + 1}\right) dx = \frac{1}{2} \sum_{k=2}^{199} \log_{10}\left(\frac{k}{k+1}\right).$$

So

$$= \frac{1}{2} \log_{10}\left(\frac{2}{3} \cdot \frac{3}{4} \cdot \frac{4}{5} \cdots \frac{199}{200}\right) = \frac{1}{2} \log_{10}\left(\frac{1}{100}\right) = -1.$$

Hence

$$\int_{100}^{1} \log_{10}\left(\frac{\lfloor 2x \rfloor}{\lfloor 2x \rfloor + 1}\right) dx = -(-1) = 1.$$

So the answer is **1**, how neat!

Question 19

Let functions $f_0(x) = |x|$ and $f_{n+1}(x) = |1 - f_n(x)|$, find

$$\int_0^{100} f_{100}(x)\,dx.$$

A 5000

B 2500

C 100

D 50

E 25

Solution 19

Answer: D

jzmaths.com

This lovely question occurred to me while I was watching R2Drew2's video on my first mock paper. In one question, he drew some very nice sketches of the sine and cosine graphs, and somehow this idea came into my head. So here it is - blame him!

The trick is to sketch successive functions $f_r(x)$ and look for a pattern. Since $|X| = |-X|$, it is easier to write the recurrence relation as

$$f_{n+1}(x) = |f_n(x) - 1|.$$

This corresponds to translating the graph down by 1 unit and then applying the modulus transformation. As you sketch the functions, you will see what is happening:



Therefore, the area described by the integral is

$$\int_0^{100} f_{100}(x)\,dx = 100 \cdot \frac{1}{2} = 50.$$

jzmaths.com

Question 20

The positive real numbers a, b, c, d form a geometric progression in this order. Given that

$$a + d = 35 \qquad \text{and} \qquad b + c = 30,$$

find the value of

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}.$$

A $\dfrac{65}{216}$

B $\dfrac{13}{7}$

C $\dfrac{13}{6}$

D $\dfrac{1}{216}$

E $\dfrac{5}{216}$

F $\dfrac{13}{210}$

jzmaths.com

Solution 20

Answer: A

Remark: This is an example of a challenging algebra problem. In problems of this kind, it is important to look for opportunities to simplify the algebra as early as possible.

Step 1: A useful identity for a 4-term geometric progression. Write $b = ar$, $c = ar^2$, $d = ar^3$ for some common ratio $r > 0$. Then

$$ad = a \cdot ar^3 = a^2r^3 \qquad \text{and} \qquad bc = ar \cdot ar^2 = a^2r^3,$$

so $ad = bc$. Calling this common product P, we have

$$\frac{1}{a} + \frac{1}{d} = \frac{a+d}{ad} = \frac{a+d}{P}, \qquad \frac{1}{b} + \frac{1}{c} = \frac{b+c}{bc} = \frac{b+c}{P}.$$

Hence

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} = \frac{(a+d)+(b+c)}{P} = \frac{65}{P}.$$

Step 2: Find $P = ad$. Using the geometric progression form,

$$a + d = a(1 + r^3) = a(1 + r)(1 - r + r^2), \qquad b + c = ar(1 + r).$$

Dividing,

$$\frac{a + d}{b + c} = \frac{1 - r + r^2}{r} = \frac{35}{30} = \frac{7}{6}.$$

So $6(1 - r + r^2) = 7r$, i.e. $6r^2 - 13r + 6 = 0$, giving $r = \frac{3}{2}$ or $r = \frac{2}{3}$.

Either choice gives the same value of $P = ad$. For $r = 3/2$: $a(1+r)(1-r+r^2) = a \cdot \frac{5}{2} \cdot \frac{7}{4} = \frac{35a}{8} = 35$, so $a = 8$. Then $d = ar^3 = 8 \cdot \frac{27}{8} = 27$, and $P = ad = 216$. For $r = 2/3$ the values of a and d swap, and P is unchanged.

Step 3: Combine.

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} = \frac{65}{216}.$$

jzmaths.com